



Bayer CropScience

Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)

**STRATEGO® Fungicide**

EPA Reg. No.: 264-779

For use on: Filberts and Triticale

Expires October 6, 2013

**Supplemental Label**

**STRATEGO® Fungicide**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read this label and the product package label before using this product. This Supplemental Label must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product package label for STRATEGO® Fungicide

<b>FILBERTS (Hazelnuts)</b>	
<b>Disease Control</b>	<b>Rate fl. oz./acre</b>
Eastern Filbert Blight ( <i>Anisogramma anomala</i> )	12-15.4
<b>Note:</b> Begin applications preventatively or when green leaf tissue becomes visible. Continue throughout the season using a 14 to 21-day spray interval. Stratego Fungicide may be applied by ground or air.	
<b>Restrictions:</b> Do not apply more than 30.4 fl.oz. of Stratego Fungicide per acre per year. Do not apply within 60 days of harvest. Do not apply more than 2 sequential applications of Stratego Fungicide before alternating with a fungicide that has a different (not Group 11) mode of action. Do not make more than 2 applications of Stratego Fungicide per acre per year. Do not graze livestock in treated areas or cut treated cover crop for feed.	

WHEAT AND TRITICALE		
Disease Control		Rate fl. oz./acre
Glume Blotch ( <i>Stagonospora nodorum</i> )	Rusts ( <i>Puccinia</i> spp.)	10.0
Leaf Blight ( <i>Septoria tritici</i> )	Tan Spot ( <i>Pyrenophora tritici-repentis</i> )	
Powdery Mildew ( <i>Erysiphe graminis</i> )		
<p><b>Note:</b> Begin applications preventively when conditions are favorable for disease development. A second application (minimum interval of 14 days) may be made if needed.</p> <p>Early season leaf disease control/suppression: Apply 4 to 8 fl. oz. /acre of Stratego Fungicide for control of early season tan spot, <i>Septoria</i>, and powdery mildew and suppression of rust.</p> <p>Stratego Fungicide may be applied by ground, air or chemigation.</p>		
<p><b>Restrictions:</b> Do not apply more than 2 applications of Stratego Fungicide per year. Do not apply after Feekes growth stage 10.5 (full head emergence). Do not apply within 35 days of harvest. Do not apply more than 2 applications of Stratego Fungicide or other Group 11 or Group 11-containing fungicide per acre per year without alternation with at least 2 applications of fungicide from a different (not Group 11) mode of action.</p> <p><b>Grazing Restrictions: (a).</b> If 1 application or a total of 10 fl. oz of Stratego Fungicide per year are applied, do not allow livestock to graze within the treated area within 30 days after application, and do not harvest the treated crop for forage within 30 days after application or for hay within 45 days after application.</p> <p><b>(b).</b> If 2 applications or a total of 20 fl. oz. of Stratego Fungicide per year are applied, do not allow livestock to graze within the treated area and do not harvest the treated crop for forage or hay.</p>		

For MEDICAL And TRANSPORTATION Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577  
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).