

SUPPLEMENTAL LABELING
For Distribution and Use Only in the State of Idaho

Sonalan[®] HFP

This supplemental label expires on December 31, 2019, and must not be used or distributed after this date.

ACTIVE INGREDIENT:	% BY WT.
Ethalfuralin: <i>N</i> -ethyl- <i>N</i> -(2-methyl-2-propenyl)-2, 6-dinitro-4-(trifluoromethyl)benzenamine.....	35.4%
OTHER INGREDIENTS:	64.6%
	TOTAL 100.0%

Contains 3 lb active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN
DANGER / PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

Oilseed – Borage, Crambe, Cuphea, Echium, Flax seed, Gold of pleasure (Camelina), Hare’s ear mustard, Lesquerella, Lunaria, Meadowfoam, Milkweed, Mustard seed, Oil radish, Poppy seed, Rapeseed (Canola Varieties Only), Sesame, and Sweet rocket (Not for use in California)

Sonalan HFP – Alone

Apply and incorporate Sonalan[®] HFP herbicide before planting. Application may be made in the fall or in the spring prior to planting canola or crambe. See instructions for fall application, application directions and soil incorporation directions on the Sonalan HFP product label.

Use the correct rate for each soil type to avoid crop injury. Prolonged wet or cold soils, deep seed placement, soil crusting, or application overlaps may combine to cause slowed or reduced emergence.

Broadcast Application Rates:

Soil Texture	Sonalan HFP (pt/acre)
coarse	1 1/2
medium	2
fine	2 1/2

Tank Mix or Overlay Recommendations

For broader spectrum weed control, other products registered for use in canola or crambe may be applied in tank mix combination or as a sequential overlay treatment following application of Sonalan HFP. Consult the manufacturer’s label for additional weeds controlled, directions for use, cautions and limitations before use. See detailed information for tank mixing in the Product Information section of this label.

USE PRECAUTIONS AND RESTRICTIONS

- Poor weed control and/or crop injury may result if directions are not followed. Uneven application, over application, or improper incorporation can result in erratic weed control or injury to the treated or rotational crops.
- Do not graze or forage crop grown in treated soil.
- Do not apply Sonalan HFP to soils which are wet, cloddy, or subject to prolonged periods of flooding because poor weed control or crop injury may result.
- Do not apply to any area not specified on the product label.
- Sonalan HFP will not control DNA (dinitroaniline) resistant goosegrass.
- **In Montana and Wyoming:** Refer to Special Rotational Crop Restrictions for the states of Montana and Wyoming in the Rotational Crop Restrictions section of the Sonalan HFP product label.
- Gold of Pleasure (Camelina) crops exhibit marginal tolerance to Sonalan HFP. Stunting or reduced stands may occur.

NET CONTENTS:

